

Evaluation of Antimicrobial Activity and Identification of Yellow Pigmented Marine Sponge-Associated Fungi from Teluk Awur, Jepara, Central Java

by Agus Trianto

Submission date: 29-Aug-2018 03:41PM (UTC+0700)

Submission ID: 994476651

File name: ponge-Associated_Fungi_from_Teluk_Awur,_Jepara,_Central_Java.pdf (1.09M)

Word count: 5756

Character count: 31283

Evaluation of Antimicrobial Activity and Identification of Yellow Pigmented Marine Sponge-Associated Fungi from Teluk Awur, Jepara, Central Java

Mada Triandala Sibero^{1,2,*}, Desy Wulan Triningsih³, Ocky Karna Radjasa^{2,4}, Agus Sabdono^{2,4}, Agus Trianto^{3,4}

¹Department of Coastal Resources Management, Faculty of Fisheries and Marine Science, Diponegoro University, Semarang, Indonesia

²Laboratory of Tropical Marine Biotechnology, Integrated Laboratory of Diponegoro University, Semarang, Central Java, Indonesia

³Laboratory of Marine Natural Product, Integrated Laboratory of Diponegoro University, Semarang, Indonesia

⁴Department of Marine Science, Faculty of Fisheries and Marine Science, Diponegoro University, Semarang, Indonesia

Abstract

Marine sponge associated fungi are known as potential source of metabolites with various biological activities. Natural pigment is one of metabolite which produced by microorganisms. Several researches reported the antimicrobial activity from natural pigment. Unfortunately there are lack of information about marine fungi natural pigment and its producer. The aims of this research were to identify yellow pigmented Indonesian marine sponge-associated fungi, to extract the pigment, and to study the antimicrobial activity of the pigment against clinical MDR bacteria and clinical pathogenic fungi. Sponge associated-fungus isolate MT23 was successfully identified as *Trichoderma parareesei*. The fungal pigment could be extracted only in methanol with yield $6.22 \pm 0.29\%$. The pigment could inhibited *Salmonella typhi* and *Escherichia coli* MDR strains. The biggest antibacterial activity was shown by concentration $1000 \mu\text{g/mL}$ against *S. typhi* with inhibition zone was 4.03 ± 0.06 mm.

Keywords: Associated fungi, pigment, *Trichoderma parareesei*

Introduction

Pigments are chemical compounds which give colours because of its ability to absorb light in the wavelength range at visible region (Delgado-Vargas *et al.*, 2000). The utilization of pigments is increasing every year due to the increase of demands in food, beverages, pharmaceutical and cosmeceutical industries (Rymbai *et al.*, 2011; Venil *et al.*, 2013). Natural pigments are known to have biological activities as its advantages over the synthetic pigments (Rostami *et al.*, 2016; Norman *et al.*, 2016; Yolmeh *et al.*, 2016).

Antimicrobial activity is one of biological activities showed by natural pigments. Salem *et al.* (2014) successfully extracted natural pigment from *Carthamus tinctorius*. The pigment displayed antimicrobial activity against Methicillin-Resistant *Staphylococcus aureus* (MRSA), *Escherichia coli* ATCC 25218 and *Candida albicans*. Another research reported that carotenoids pigment were able to inhibit *Salmonella enteritidis* (PTCC 1709) and *E. coli* (PTCC1260) (Rosli *et al.*, 2016). In addition, crude extract of pigment from *Streptomyces* sp. D25 was reported to have antibacterial activity against biofilm forming bacteria such as *Pseudomonas* sp. P1, *Bacillus* sp. P13 and *Alcaligenes* sp. M28 (Radhakrishnan *et al.*, 2016).

Marine microorganisms are known as potential producer of natural pigment with great antibacterial activity. There are several microorganisms are known life associated

*Corresponding author:

Mada Triandala Sibero
Faculty of Fisheries and Marine Science,
Diponegoro University, Jalan Imam Bardjo, SH.,
Semarang 50241, Indonesia
E-mail: madatriandala@hotmail.com

with marine organisms especially sponges such as bacteria, actinomycetes and fungi (Idraningrat *et al.*, 2016; Wu *et al.*, 2016). Ibrahim *et al.* (2014) studied antimicrobial activity of prodigiosin (red pigment) from sponge-associated bacteria *Serratia marcescens* IBRL USM 84. This pigment greatly inhibited MRSA and had good activity inhibited *Bacillus cereus*, *Bacillus licheniformis* and *Agrobacterium* *13* *tafaciens*. In 2015, Suresh *et al.* (2015) observed antibacterial activity of red pigment produced by *Halolactibacillus alkaliphilus* MSRDI which isolated from seaweed against *S. aureus* and *Salmonella typhi*. In addition, Srilekha *et al.* (2016) successfully isolated marine pigmented bacteria and extracted the pigment then studied its antimicrobial activity. Result showed the pigment had great antimicrobial activity against bacteria such as *S. aureus*, *Proteus vulgaris*, *K. pneumoniae*, *E. coli*, and fungi such as *Fusarium sp.*, *C. albicans*, *Mucor sp.*, and *A. flavus* with inhibition zone range 18–28 mm. These reports show great potential of natural pigments from marine bacteria as antibacterial agent.

However, there are still a few researches about the potential of natural pigments from marine fungi and its antibacterial activity against clinical multidrug resistant bacteria (MDR) and clinical pathogenic fungi. MDR bacteria are bacteria which already resistant to several antibiotics that should be effective to inhibit the growth of the bacteria (Magiorakos *et al.*, 2012). This is a big issue for public health. The aims of this research were to identify Indonesian yellow pigmented sponge-associated fungi, to extract the pigment, and to study the antimicrobial activity of the pigment against clinical MDR bacteria and clinical pathogenic fungi.

Materials and Methods

Media and chemicals

This research used Malt Extract Agar (MEA) M137 base with mycological 8-*septone* and MacConkey M081B agar from HiMedia Laboratories Pvt. Ltd. (Swastik Disha Business Park, Via Vadhani, Ind. Est., LBS Marg, Mumbai-400086, India), Nutrient Agar CM0003

from Oxoid Ltd. (Basingstoke, Hampshire, England), Mueller Hinton Agar (MHA) ACC, to CLSI 1.05435.0500 from Merck KGaA (Darmstadt, Germany) while agar 36 was bought from Promega. Solvents such as methanol, ethyl acetate and n-hexane were bought from Merck. DNA extraction used Chelex®100 (Sigma-Aldrich), PCR master mix GoTaq Green Master Mix from Promega Corporation (2800 Woods Hollow Road Madison, WI 53711-5399 U.S.A) while 35 primer (ITS1 and ITS4) from Macrogen (1330 Piccard Drive Rockville, MD, 20850 United States).

Fungal isolate

Fungus MT23 was culture collection of Marine Fungi Division of Tropical Marine Biotechnology Laboratory, Integrated Laboratory of Diponegoro University, Semarang, Central Java, Indonesia. This fungus was isolated from unidentified marine sponge from Teluk Awur Bay, Jepara, Central Java.

MDR bacteria and clinical pathogenic fungi isolates

MDR bacteria and clinical pathogenic fungi were obtained from Dr. Kariadi General Hospital Medical Center, Semarang, Central Java, Indonesia and Diponegoro National Hospital, Semarang, Central Java, Indonesia. Bacteria *E. coli*, *S. typhi* and fungi *Malassezia furfur* and *Trichophyton rubrum* were used in this study.

Fungal culture

Fungus was recultured according to Sibero *et al.* (2016) with particular modifications. Approximately 1 cm² fungus was taken from 34 stock culture then transferred to fresh Malt Extract Agar (MEA) and incubated in room temperature (23°C) for 7 days with addition of chloramphenicol 5% (%w/w). After that, the morphology of the recultured fungus was compared to the culture stock. The same morphology indicate the fungus was not contaminated. Then fungus MT23 was reculture to new fresh MEA without chloramphenicol, pH 5.4, illumination using LED lamp (Philip LED 20 watt) and incubated in room temperature (23°C) for 14 days.

MDR bacteria and clinical pathogenic fungi

Gram-negative bacteria *E. coli* and *S. typhi* were revived on MacConkey Agar while *M. furfur* and *T. rubrum* were revived on Potato Dextrose Agar (PDA). The bacteria were incubated for 24 h in 32°C while the fungi were incubated in at 27°C for 24 h.

Fungal identification

Morphological approach

Fungus MT23 was recultured on MEA for 5 days with slide cultured method (Qiu *et al.* 2005). Sterilized MEA was cut with size 2 cm² then transferred and put onto object glass then covered by cover glass. After that, fungal mycelia were inoculated on each side of the MEA then incubated at room temperature (set at 23°C). After 5 days of incubation, object and cover glasses were covered by fungal mycelia. MEA was separated from the glasses. Lactophenol cotton blue was dropped on the object and cover glasses then covered by its pairs. After that, the morphology of the fungi was observed under microscope.

Molecular approach

DNA of fungus MT23 was extracted using Chelex method from Turan *et al.* (2015) with particular modifications. Seven days old of mycelium were immersed in 100 µL ddH₂O and 1000 µL of 0.5% saponin for 24 h in chilling temperature (4°C). Then it was separated by centrifugator with 12000 rpm for 10 min. The supernatant was discarded while template was added 100 µL of ddH₂O followed by addition of 50 µL of 20% Chelex 100. Then it was heated in water bath at 80°C for 5 min. After that, the mixture was homogenized using vortex for 10 s and reheated for 5 min. After reheated, the DNA in mixture were separated using centrifugation at 12000 rpm for 10 min. Then the supernatant were transferred to new microtube and stored at -20°C for 24h.

DNA of fungus MT23 was amplified based on Internal Transcribed Spacer (ITS) region. The amplification using primer 1 µL of ITS 1 (5'-TCC GTA GGT GAA CCT GCG G-3') and 1 µL of ITS 4 (5'-TCC TCC GCT TAT TGA TAT GC-3'). The primers were mixed

with 12.5 µL of GoTaq Green Master Mix Promega, 0.5 µL of DNA extract and 10 µL of ddH₂O. The PCR profile consists of 3 min of preheat at 93°C for 3 min, 30 cycles for denaturation at 95°C (1 min), annealing at 56.1°C (1 min) and extension at 72°C (1 min). Final extension was done at 72°C for 7 min. The quality of PCR product was checked using gel electrophoresis in agarose 1%. Then PCR product was sent to 1st Base Laboratories Sdn Bhd, Malaysia for DNA sequencing. DNA sequence was analyzed to its homology using Basic Local Alignment Search Tool (www.ncbi.nlm.nih.gov). Phylogenetic tree was reconstructed using MEGA 7.0 software package (Tamura *et al.* 2011). Neighbor-joining was applied as the statistical method with 1000 bootstrap replications.

Pigment extraction

Yellow pigment from fungus MT23 was extracted using solid-liquid extraction (Manikkam *et al.* 2015). The mycelia were separated from media, then the media were cut into smaller size and dried using silica gel in desiccator for 48 h. After that, solvent optimization for pigment extraction was done by immersing the dried agar in methanol, ethyl acetate and n-hexane with agitation using a shaker (118 rpm) for 24 h then filtered using filter paper (Macherey-Nagel 640d Ø 1125 mm). The filtrate was evaporated using rotary evaporator (Eyela SB-1100) set at 38°C. The pigment yield percentage was calculated using following equation:

$$\% \text{ Pigment yield} = \frac{\text{Pigment yield (gr)}}{\text{Agar weight (gr)}} \times 100\%$$

Determination of inhibition zone

Antibacterial activity

Antibacterial activity of the pigment was evaluated using agar-disc diffusion method from Balouiri *et al.* (2016) with several modifications. Pigment extract was dissolved in methanol and diluted to five concentrations (50, 125, 250, 500 and 100 µg/mL). *S. typhi* and *E. coli* strain MDR were grown on MacConkey Agar for 24 h (32°C) then diluted in physiological saline solution into 0.5

McFarland. Then the bacterial solution was inoculated to MHA using sterile cotton swab. Each extract (30 µl) was injected onto sterilized paper disc then placed on the MHA and incubated at 32°C for 24 h. Amoxicillin + Clavulanic acid (30 µg/disc) was used as positive control. The presence of clear zone indicated the antibacterial activity of the pigment extract.

Antifungal activity

This assay was done according to Bhalodia and Shukla (2011) and Balouiri *et al.* (2016) with modifications. Pigment extract was dissolved in methanol and diluted to five concentrations (50, 125, 250, 500 and 100 µg/mL). *M. furfur* and *T. rubrum* were grown on PDA for 24 h at 27°C. Then the fungi were inoculated using pour plate method. Each extract (10 µl) was injected onto sterilized paper disc then placed on the PDA and incubated at 27°C for 24 h. Nystatin was used as positive control in this study. The presence of clear zone on PDA indicated the antifungal activity of the pigment extract.

Data analysis

The impact of solvent to pigment yields was calculated with One Way Analysis of Variance (ANOVA) with $P < 0.05$ while antibacterial and antifungal activities were calculated using Two Way ANOVA with $P < 0.05$. The data was analyzed using SPSS 5.0.

Results and Discussion

Fungal identification

Fungus MT23 is a marine sponge associated fungi which is isolated from sponge from Teluk Awur, Jepara, Central Java. Macroscopic characteristics of fungus MT23 were white colour mycelium with green radial growth ring, hyphae concentrated and produced yellow colour in the media. This fungus grown well in room temperature (23°C) and changed the colour of MEA from slightly orange to yellow colour. Colour change was started at fourth day of cultivation and successfully changed the colour of whole media at day 14 (Figure 1). Physical conditions of environment including pH and temperature

were reported to give impact to the fungal pigment production. A fungus *Penicillium purpurogenum* GH2 produced highest yield of red pigment after 240 h (10 days) of cultivation with pH 5 at 24°C (Mendéz *et al.* 2011). On the other hand, Velmurungan *et al.* (2010) induced fungal pigment production using colour of lights. They figured out that the darker lights like red (780–622 nm) and blue (492–455) gave higher pigment productions compared to green (577–492 nm), yellow (597–577 nm) and white lights in 5 different fungi. This phenomenon could be explained by the postulating of the existence of photoreceptor in fungi. However, there was no clear explanation about the mechanisms of the light to induce pigment productions in fungi. Furthermore, Blumenstein *et al.* (2005) suggested that a phytochrome type of system may be operative in particular fungi, in his case was *A. tulans*. The phytochrome suggested able to sense the red and far-red light through photointerconversion between the two stable conformations.

Figure 2 shows the morphology of fungus MT23 under microscope. This fungus had hyaline and thin mycelium with septum. The conidiophore were branching with lageniform phialide with green conidia, smooth, eguttulate, and ellipsoidal to cylindrical form. According to its characteristics, fungus MT23 was judged as member of genus *Choderma* (Atanasova *et al.*, 2010; Rahman *et al.*, 2011; Qin and Zhuang, 2016). To confirm the accuracy of morphological identification, we did identification through molecular approach.

Fungal DNA was extracted using Chelex 100 method (Turan *et al.*, 2015). This method is commonly used to extract microorganisms' DNA (Liu *et al.*, 2015; Susilowati *et al.*, 2015). Chelex 100 able to prevent DNA damage by inhibiting DNase activity (Walsh *et al.*, 1997). Fungal mycelium was immersed in saponins and ddH₂O for overnight. Saponins is detergent agent to lyse the cells (Dong *et al.*, 1997). To amplify fungal DNA, ITS1 and ITS4 were used as primer. These primers will amplify DNA from end of 18S rRNA gene region, internal transcribed spacer 1, 5.8S rRNA gene

region, internal transcribed spacer 2 region and the beginning of 28S rRNA gene region (White *et al.*, 1990; Larena *et al.*, 1999; Anderson *et al.*, 2003; Bellemain *et al.*, 2010; Schoch *et al.*, 2012). Figure 3 shows the location which amplified by ITS primer.




Figure 1. a) 4 days old of mycelium, b) 4 days old of reverse side, c) 14 days old of reverse side.




Figure 2. a) conidiophore, b) mycelium, c) conidia.

Fungal DNA amplification was done using polymerase chain reaction (PCR) the quality of PCR product was checked using gel electrophoresis. The result of visualization of fungal DNA quality is shown by Figure 4.




Figure 3. Diagram of the rDNA region of fungi and ITS primer (Source: Bellemain *et al.*, 2010).

Figure 4 shows that the fungal DNA was successfully amplified in region between 500 to 1000 bp. This result is supported by the statement of Nilsson *et al.* (2009) who stated that the fungal DNA is normally obtainable in a single round of Sanger DNA sequencing in 650 bp region. PCR product was sequenced to get the sequence of fungal

DNA, and then the sequence was compared to that of the database in gene bank using BLAST homology. According to fungal DNA comparison, fungus MT23 was 99% similar to *Trichoderma parareesei* ATCC MYA-4777 (Houseknecht *et al.*, 2011). Phylogenetic tree of fungus MT23 is shown by Figure 5.




Figure 4. Visualization of DNA band of fungus MT23.

Result of morphology observation was supposed fungus MT23 was member of genus *Trichoderma* and supported by the molecular identification which successfully identified fungus MT23 as *T. parareesei*. Genus *Trichoderma* is well known as terrestrial fungus and give advantages for agricultural sector (Rinu *et al.*, 2013; Saravanakumar *et al.*, 2013; El Komy *et al.*, 2015; Hamed *et al.*, 2015). The finding of *T. parareesei* as marine sponge associated fungi showed that this fungus was a marine facultative (marine derived) fungi. Several species from *Trichoderma* such as *T. longibrachiatum*, *T. harzianum*, and *T. atroviride* were reported to tolerate to the increasing of salinity (Gal-Hemed *et al.* 2011). Genus of *Trichoderma* was also known to produced pigment and bioactive compounds (Rubeena *et al.*, 2013; Vacondio *et al.*, 2015; Benkada *et al.*, 2016).

Pigment extraction

Fungus MT23 changed the colour of MEA to yellow because it produced extracellular pigment. Extracellular pigments production is indicated by colour changing of the environment where the fungi live. Previous research successfully identified black extracellular pigment which produced by endophytic fungi as melanin having photoprotector activity (Sibero *et al.*, 2016). Fungus MT23 produced yellow pigment started at the fourth day of cultivation and




Figure 5. Phylogenetic tree of fungus MT23.

ready to be harvested on day fourteenth. Pradeep and Pradeep (2013) studied the optimum cultivation condition for pigment production of fungus *Fusarium moniliforme*. The fungus had highest pigment absorbance (AU 500^{nm}) at temperature 28–30°C, pH 5.5, glucose as carbon source, peptone as nitrogen source and cultivated for 8 days. Yellow pigment from fungus MT23 were extracted using organic solvents. Environmental conditions have important role in fungi growth. Pradeep and Pradeep (2013) studied the optimum cultivation condition for growth of fungus *Fusarium moniliforme*. The fungus had highest biomass (g_g) at temperature 28°C, pH 5.5, glucose as carbon source, peptone as nitrogen source, KH₂PO₄ as source of salt and addition of methionine as additional amino acid. Stationary phase of this fungus was on eighth day of cultivation. In this study, we used Malt Extract Agar (MEA) from HiMedia M173 which contained malt extract and mycological peptone. Peptide and amino acid in peptone are easily metabolized by fungi which induced the production of fungal metabolites, including pigment (Calestino *et al.*, 2014). Da Costa Souza *et al.* (2016) used several media including malt extract to induced pigment production in filamentous fungi. As a result, pigment production in malt extract was higher than other medium. In addition,

nutrient content in media can regulate the expression of genes to activate the metabolic path way for pigments production (Pradeep *et al.*, 2013).

Organic solvents such as n-hexane, ethyl acetate, chloroform, methanol and dichloromethane were usually used to extract natural pigments from microorganisms (Saravanan and Radhakrishnan, 2016; Weber *et al.*, 2016). In this research, n-hexane, ethyl acetate and methanol were used to extract fungal pigment. The purpose of this step was to know the best solvent to obtain highest fungal pigment yield. The result of pigment extraction is shown by Figure 6 and Table 1.

Table 1 shows that only methanol was able to extract fungal extracellular pigment from fungus MT23. Statistic analysis showed that solvents gave significant difference to the fungal pigment yield ($P < 0.05$). The polarity of pigment will influence the suitability of



Figure 6. The result of maceration of agar media in organic solvents: a) in ethyl acetate, b) in n-hexane, c) in methanol.

Table 1. Fungal pigment yield with different solvents.

Solvent	Fungal pigment yield (%)
n-Hexane	0
Ethyl acetate	0
Methanol	6.26±0.29

optimum solvent to extract the pigment. Several pigments such as astaxanthin, chlorophyll a, chlorophyll b and xanthophyll were able to be extracted with methanol (Sasidharan *et al.*, 2013, Sumanta *et al.*, 2014; Bhat and Marar 2015).

Antibacterial and antifungal activities

Fungal pigments are known to produce antimicrobial activity. Geweely (2011) successfully investigated antimicrobial of several fungal pigments. The result showed that fungus *Aspergillus nidulans*, *Fusarium moniliforme*, *Penicillium purpurogenum*, and *Phoma herbarum* had weak to moderate antimicrobial activity against dermatophyte fungi, non dermatophyte fungi and several bacteria. Another research showed antibacterial activity of astaxanthin pigment from marine yeast (Ushakumari and Ramanujan, 2013). Fungal pigment from fungus MT23 was tested against several clinical pathogenic microorganisms. The result of antibacterial and antifungal is shown in Table 2.

According to Table 2, fungal pigment from MT23 showed weak antibacterial activity against clinical MDR bacteria *S. typhi* and *E. coli*. The biggest antibacterial activity showed at the concentration of 1000 µg/mL against *S. typhi*. The pigment didn't show any antifungal activity against clinical pathogenic fungi. The

weak antimicrobial activity means that this fungal pigment is not potential as a source of antibiotic against clinical pathogenic and MDR microorganisms. Carotenoid is one of natural pigment which produced by fungi with yellow, orange to reddish color. The production of this pigment is related to stress tolerance or with the synthesis of physiologically active by product (Avalos and Limón, 2015). Carotenoid pigment is mostly reported has antimicrobial and antioxidant potential (Ernawita *et al.*, 2016; Yoo *et al.*, 2011). The antimicrobial mechanisms of this pigment could lead to the accumulation of lysozyme enzyme that digest bacterial cell walls (Abu-Ghannam *et al.*, 2013).

Conclusions

Sponge associated-fungus isolate MT23 was successfully identified using morphological and molecular approaches as member of *Trichoderma* and had 99% similarity to *Trichoderma parareesei*. The fungal pigment could be extracted only in methanol with yield 6,22±0,29%. The pigment showed weak antibacterial activity against *S. typhi* and *E. coli* strain MDR, furthermore the pigment did not show any activity against clinical pathogenic fungi.

Acknowledgements

We thank Directorate of Research and Community Development, Ministry of Research, Technology and Higher Education for the research grant through Program Magister Menuju Doktor Bagi Sarjana Unggulan (PMDSU) scheme, funding in the year 2016, No: 2993-12/UN7.5.1/PG/2016.

Table 2. Antibacterial and antifungal activities of fungal pigment MT23.

Concentration (µg/mL)	Diameter of Inhibition Zone (mm)			
	Clinical MDR Bacteria		Clinical Pathogenic Fungi	
	<i>S. typhi</i>	<i>E. coli</i>	<i>M. furfur</i>	<i>T. rubrum</i>
50	0.00	0.00	0.00	0.00
125	0.00	0.00	0.00	0.00
250	0.00	0.00	0.00	0.00
500	1.13±1.00 ^a	0.00	0.00	0.00
1000	4.03±0.06 ^b	2.50±0.01 ^a	0.00	0.00
Positive control	6.40±0.24 ^c	11.58±0.00 ^b	19.70±0.38	20.10±0.02
Negative control	0.00	0.00	0.00	0.00

Superscript shows statistically significant difference ($P < 0.05$)

References

- Abu-Ghannam, N., and Rajauria, G. (2013). Antimicrobial activity of compounds isolated from algae. In *Functional Ingredients from Algae for Foods and Nutraceuticals*; Domínguez, H., Ed. Cambridge (UK): Woodhead Publishing.
- Anderson, I.C., Campbell, C.D., Prosser, J.I. 2003. Potential bias of fungal 18S rDNA and internal transcribed spacer polymerase chain reaction primers for estimating fungal biodiversity in soil. *Environ. Microbiol.*, 5(1), 36-47.
- Atanasova, L., Jaklitsch, W.M., Komoń-Zelazowska, M., Kubicek, C.P., Druzhinina, I.S. 2010. Clona species *Trichoderma parareesei* sp. nov. likely resembles the ancestor of the cellulase producer *Hypocrea jecorina*/ *T. reesi*. *Appl. Environ. Microbiol.*, 76(21), 7259-7267.
- Avalos, J., Limon, M.C. 2015. Biological roles of fungal carotenoids. *Curr. Genet.*, 61(3), 309-324.
- Balouiri, M., Sadiki, M., Ibensouda, S.K. 2016. Methods for in vitro evaluating antimicrobial activity: A review. *J. Pharm. Anal.*, 6, 71-79.
- Bellemain, E., Carlsen, T., Brochmann, C., Coissac, E., Taberlet, P., Kauserud, H. 2010. ITS as an environmental DNA barcode for fungi: an in silico approach reveals potential PCR biases. *BMC Microbiol.*, 10(89), 1-9.
- Benkada, M.M., Pouchus, Y.F., Vêrite, P., Pagniez, F., Caroff, N., Ruiz, N. 2016. Identification and biological activities of long-chain peptaibols produced by a marine-derived strain of *Trichoderma longibrachiatum*. *Chem. Biodiver.*, 13, 521-530.
- Bhalodia, N.R., and Shukla, V.J. 2011. Antibacterial and antifungal activities from leaf extracts of *Cassia fistula* L.: An ethnomedicinal plant. *J. Adv. Pharm. Technol.*, 2(2), 104-109.
- Bhat, M.R., and Marar, T. 2015. Media optimization, extraction and partial characterization of an orange pigment from *Salinicoccus* sp. MKJ 997975. *Int. J. Life. Sci. Biotech. Pharm. Res.*, 4(2), 85-89.
- Blumenstein, A., Vienken, K., Purschwitz, J., Tasler, R., Frankenberg, N., Fischer, R. 2005. The phytochrome FphA controls development in the filamentous fungus *Aspergillus nidulans*. *Curr. Biol.*, 15, 1833-1838.
- Celestino, J., dos, R., de Carvalho, L.E., Lima, M. da P., Lima, A.M., Ogusku, M.M., de Souza, J.V.B. 2014. Bioprospecting of Amazon soil fungi with the potential for pigment production. *Process Biochem.*, 49, 569-575.
- de Costa Souza, P.N., Grigoletto, T.L.B., de Moraes, L.A.B., Abreu, L.M., Guimaraes, L.H.S., Santos, C., Galvao, L.R., Cardoso, P.G. 2016. Production and chemical characterization of pigments in filamentous fungi. *Microbiol.*, 162, 12-22.
- Delgado-Vargas, F., Jiménez, A.R., Paredes-López, O. 2000. Natural pigments: carotenoids, anthocyanins, and betalains-characteristics, biosynthesis, processing, and stability. *Crit. Rev. Food. Sci.*, 40(3), 173-289.
- Dong, Z., Saikumar, P., Weinberg, J.M., Venkatachalam, M.A. 1997. Internucleosomal DNA cleavage triggered by plasma membrane damage during necrotic cell death. Involvement of serine but not cysteine proteases. *Am. J. Pathol.*, 151(5), 1205-1213.
- El Komy, M.H., Saleh, A.A., Eranthodi, A., Molan, Y.Y. 2015. Characterization of novel *Trichoderma asperellum* isolates to selects effective biocontrol agents against tomato *Fusarium* wilt. *Plant Pathol. J.*, 31(1), 50-60.
- Ernawita, Wahyuono, R.A., Hesse, J., Hippler, U.C., Elsner, P., Böhm, V. 2016. Carotenoids of indigenous citrus species from Aceh and its in vitro antioxidant, antidiabetic and antibacterial activities. *Eur. Food. Res. Technol.*, 242(11), 1869-1881.
- Gal-Hemed, I., Atanasova, L., Komoń-Zelazowska, M., Druzhinina, I.S., Viterbo, A., Yarden, O. Marine isolates of

- Trichoderma* spp. as potential halotolerant agents of biological control for arid-zone agriculture. *App. Environ. Microbiol.*, 77(15), 5100-5109.
- Geweely, N.S. 2012. Investigation of the optimum condition and antimicrobial activities of pigments from four potent pigment-producing fungal species. *J. Life Sci.*, 5, 697-711.
- Hamed, R.E., Awad, H.M., Ghazi, E.A., El-Gamal, N.G., Shehata, H.S. 2015. *Trichoderma asperellum* isolated from salinity soil using rice straw waste as biocontrol agent for cowpea plant pathogens. *J. App. Pharm. Sci.*, 5(2), 91-98.
- Houseknecht, J.L., Suh, S.O., Zhou J. 2011. *Trichoderma parareesei* ATCC MYA-4777 ITS region; from TYPE material. [https://www.ncbi.nlm.nih.gov/nucleotide/1051341416?report=genbank&log\\$=nucalign&blast_rank=2&RID=5RZZ5WVG01R](https://www.ncbi.nlm.nih.gov/nucleotide/1051341416?report=genbank&log$=nucalign&blast_rank=2&RID=5RZZ5WVG01R) (Accessed on 22 Januari 2016).
- Ibrahim, D, Nazari, T.F., Kassim, J, Lim S.H. 2014. Prodigiosin-an antibacterial red pigment produced by *Serratia marcescens* IBRL USM 84 associated with a marine sponge *Xestospongia testudinaria*. *J. App. Pharm. Sci.*, 4(10), 001-006.
- Idraningrat, A.A.G., Smidt, H., Sipkema, D. 2016. Bioprospecting sponge-associated microbes for antimicrobial compounds. *Mar. Drugs*, 14(5), 87.
- Larena, I., Salazar, O., González, V., Julián, M.C., Rubio, V. 1999. Design of a primer for ribosomal DNA internal transcribed spacer with enhanced specificity for Ascomycetes. *J. Biotechnol.*, 75, 187-194.
- Liu, L., Wang, C.L., Peng, W.Y., Yang, J., Lan, M.Q., Zhang, B., Li, J. B., Zhu, Y.Y., Li, C.Y. 2015. Direct DNA extraction method on an obligate parasitic fungus from infected plant tissue. *Genet. Mol. Res.*, 14(4), 18546-18551.
- Magiorakos, A.P., Srinivasan, A., Carey, R.B., Carmeli, Y., Falagas, M.E., Giske, C.G., Harbarth, S., Hindler, J.F., Kahlmeter, G., Olsson-Liljequist, B., Paterson, D.L., Rice, L.B., Stelling, J., Struelens, M.J., Vatopoulos, A., Weber, J.T., Monnet, D.L. 2012. Multidrug-resistant, extensively drug-resistant and pandrug-resistant bacteria: an international expert proposal for interim standard definitions for acquired resistance. *Clin. Microbiol. Infect.*, 18(3), 268-281.
- Manikkam, R., Venugopal, G., Ramasamy, B., Kumar, V. 2015. Effect of critical medium components and culture conditions on antitubercular pigment production from novel *Streptomyces* sp. D25 isolated from Thar desert, Rajasthan. *J. App. Pharm. Sci.*, 5(06), 015-019.
- Méndez, A., Pérez, C., Montanez, J. C., Martínéz, G., Aguilar, C. N. 2011. Red pigment production by *Penicillium purpurogenum* GH2 is influenced by pH and temperature. *J. Zhejiang Univ.*, 12(12), 961-968.
- Nilsson, R.H., Ryberg, M., Abarenkov, K., Sjökvist, E., Kristiansson, E. 2009. The ITS region as a target for characterization of fungal communities using emerging sequencing technologies. *FEMS Microbiol. Lett.*, 296, 97-101.
- Norman, M., Bartczak, P., Zdarta, J., Ehrlich, H., Jesionowski. 2016. Anthocyanin dye conjugated with *Hippospongia communis* marine demosponge skeleton and its antiradical activity. *Dyes Pigm.*, 134, 541-552.
- Pradeep, F.S. and Pradeep, B.V. 2013. Optimization of pigment and biomass production from *Fusarium moniliforme* under submerged fermentation conditions. *Int. J. Pharm. Pharm. Sci.*, 5(3), 526-535.
- Pradeep, F.S., Begam, M.S., Palaniswamy, M., Pradeep, B.V. 2013. Influence of culture media on growth and pigment production by *Fusarium moniliforme* KUMBF1201 isolated from paddy field soil. *World Appl. Sci. J.*, 22, 70-77.
- Qin, W.T. and Zhuang, W.Y. 2016. Seven wood-inhabiting new species of the genus *Trichoderma* (Fungi, Ascomycota) in Viride clade. *Sci. Rep.*, 6(27074), 1-14.
- Qiu W.Y., Yao, Y.F., Zhu, Y.Y., Zhang, Y.M., Zhou P., Jin, Y.Q., Zhang, B. 2005. Fungal Spectrum Identified by a New

- Slide Culture and In Vitro Drug susceptibility Using Etest in Fungal Keratitis. *Curr. Eye. Res.*, 30, 1113-1120.
- Radhakrishnan, M., Gopikrishnan, V., Vijayalakshmi, G., Kumar, V. 2016. In vitro antioxidant activity and antimicrobial activity against biofilm forming bacteria by the pigment from desert soil *Streptomyces* sp D25. *J. Appl. Pharm. Sci.*, 6(6), 148-150.
- Rahman, A., Begum, M.F., Rahman, M., Bari, M.A., Illas, G.N.M., Alam, M.F. 2011. Isolation and identification of *Trichoderma* species from different habitats and their use for bioconversion of solid waste. *Turk. J. Biol.*, 35, 183-194.
- Rinu, K., Sati, P., Pandey, A. 2013. *Trichoderma gamsii* (NFFCCI 2177): A newly isolated endophytic, psychrotolerant, plant growth promoting, and antagonistic fungal strain. *J. Basic. Microbiol.*, 54(5), 408-417.
- Rostami, H., Hamed, H., Yolmeh, M. 2016. Some biological activities of pigments extracted from *Micrococcus roseus* (PTCC 1411) and *Rhodotorula glutinis* (PTCC 5257). *Int. J. Immunopathol. Pharmacol.*, 29(4), 684-695.
- Rubeena, M., Neethu, K., Sajith, S., Sreedevi, S., Priji, P., Unni, K.N., Josh, M.K.S., Jisha, V.N., Pradeep, S., Benjamin, S. 2013. Lignocellulotic activities of novel strain of *Trichoderma harzianum*. *Adv. Biosci. Biotechnol.*, 3, 214-221.
- Rymbai, H., Sharma, R.R., Srivastav, M. 2011. Bicolorants and its implications in health and food industry-a review. *Int. J. PharmTech. Res.*, 3(4), 2228-2244.
- Salem, N., Msaada, K., Elkahoui, S., Mangano, G., Azaeiz, S., Slimen, I.B., Kefi, S., Pintore, G., Limam, F., Marzouk, B. 2014. Evaluation of antibacterial, antifungal, and antioxidant activities of Safflower natural dyes during flowering. *BioMed. Res. Int.*, 2014, 1-10.
- Saravanakumar, K., Arasu, V.S., Kathiresan, K. 2013. Effect of *Trichoderma* on soil phosphate solubilization and growth improvement of *Avicennia marina*. *Aquat. Bot.*, 104, 101-105.
- Saravanan, D., and Radhakrishnan, M. 2016. Antimicrobial activity of pigments produced by fungi from Western Ghats. *J. Chem. Pharm. Res.*, 8(1), 634-638.
- Sasidharan, P., Raja, R., Karthik, C., Sharma, R., Arulselvi, P. I. 2013. Isolation and characterization of yellow pigmented producing *Exiguobacterium* sps. *J. Biochem. Toxicol.*, 4(4), 632-635.
- Schoch, C.L., Seifert, K.A., Huhndorf, S., Robert, V., Spouge, J.L., Levesque, C. A., Chen, W., Concorium, F.B. 2012. Nuclear ribosomal internal transcribed spacer (ITS) region as a universal DNA barcode marker for fungi. *Proc. Natl. Acad. Sci.*, 109(16), 6241-6246.
- Sibero, M.T., Tarman, K., Hanif, N. 2016. Characterization and photoprotector activity of endophytic fungal pigments from coastal plant sarang semut (*Hydnophytum formicarum*). *J. Pengolah. Hasil. Perikan. Indones.*, 19(1), 1-8.
- Srilekha, V., Krishna, G., Srinivas, S.V., Charya, S.M.A. 2016. Isolation and screening of marine pigmented bacteria from Nellore Coast for antimicrobial studies. *J. Cell. Tissue. Res.*, 16(1), 5413-5419.
- Sumanta, N., Haque, C.I., Nishika, J., Suprakash, R. 2014. Spectrophotometric analysis of chlorophylls and carotenoids from commonly grown fern species by using various extracting solvents. *Res. J. Chem. Sci.*, 4(9), 63-69.
- Suresh, M., Renugadevi, B., Brammavidhya, S., Iyapparaj, P., Anantharaman, P. 2015. Antibacterial activity of red pigment produced by *Halolactibacillus alkaliphilus* MSR1—an isolate from seaweed. *Appl. Biochem. Biotechnol.*, 176(1), 185-195.
- Susilowati, R., Sabdono, A., Widowati, I. 2015. Isolation and characterization of bacteria associated with brown algae *Sargassum* spp. from Panjang Island and their antibacterial activity. *Procedia Environ. Sci.*, 23, 240-246.
- Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M., Kumar, S. 2011. MEGA5: Molecular evolutionary genetics

- analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol. Biol. Evol.*, 28(10), 2731-2739.
- Turan, C., Nanni, I.M., Brunelli, A., Collina, M. 2015. New rapid DNA extraction method with Chelex from *Venturia inaequalis* spores. *J. Microbiol. Methods*, 115, 139-143.
- Ushakumari, U.N.U., and Ramanujan, R. 2013. Isolation of astaxanthin from marine yeast and study of its pharmacological activity. *Int. Curr. Pharm. J.*, 2(3), 67-69.
- Vacondio, B., Birolli, W., Ferreira, I., Porto, A.L.M. 2015. Biodegradation of pentachlorophenol by marine-derived fungus *Trichoderma harzianum* CBMAI 1677 isolated from ascidian *Didemnum ligulum*. *Biocatal. Agric. Biotechnol.*, 4(2), 266-275.
- Velmurungan, P., Lee, Y.H., Venil, C.K., Laksmanaperumalsamy, P., Chae, J.C., Oh, B.T. 2010. Effect of light on growth, intracellular and extracellular pigment production by five pigment-producing filamentous fungi in synthetic medium. *J. Biosci. Bioeng.*, 109(4), 345-350.
- Venil, C.K., Zakaria, Z.A., Ahmad, W.A. 2013. Bacterial pigments and their application. *Process Biochem.*, 48, 1065-1079.
- Walsh, P.S., Metzger, D.A., Higuchi, R. 2013. Chelex 100 as a medium for simple extraction of DNA for PCR-based typing from forensic material. *BioTechniques*, 54(3), 134-139.
- Weber, G.L., Boonloed, A., Naas, K.M., Koesdjojo, M.T., Remcho, V.T., Robinson, S.C. 2016. A method to stimulate production of extracellular pigments from wood-degrading fungi using a water carrier. *Curr. Res. Environ. Appl. Mycol.*, 6(3), 218-230.
- White, T.J., Bruns, T., Lee, S., Talyor, J. (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis, M.A., Gelfand, D.H., Sninsky, J.J. and White, T. J. Eds. *PCR protocols: a guide to methods and applications*. San Diego (US): Academic Press. pp. 315-322.
- Wu, B., Oesker, V., Wiese, J., Schmaljohann, R., Imhoff, J. F. 2014. Two new antibiotic pyridones produced by a marine fungus *Trichoderma* sp. strain MF106. *Mar. Drugs*, 12, 1208-1219.
- Yolmeh, M., Hamed, H., Khomeiri, M. 2016. Antimicrobial activity of pigments extracted from *Rhodotorula glutinis* against some bacteria and fungi. *Zahedan J. Res. Med. Sci.*, 18(12), 1-5.
- Yoo, A.Y., Alnaeeli, M., Park, J.K. 2016. Production control and characterization of antibacterial carotenoids from the yeast *Rhodotorula mucilaginosa* AY-01. *Process Biochem.*, 51(4), 463-473.

Evaluation of Antimicrobial Activity and Identification of Yellow Pigmented Marine Sponge-Associated Fungi from Teluk Awur, Jepara, Central Java

ORIGINALITY REPORT

15%

SIMILARITY INDEX

%

INTERNET SOURCES

15%

PUBLICATIONS

%

STUDENT PAPERS

PRIMARY SOURCES

1

"Plant-Microbe Interactions in Agro-Ecological Perspectives", Springer Nature, 2017

Publication

1%

2

Palanivel Velmurugan, Yong Hoon Lee, Chidambaram Kulandaisamy Venil, Perumalsamy Lakshmanaperumalsamy et al. "Effect of light on growth, intracellular and extracellular pigment production by five pigment-producing filamentous fungi in synthetic medium", Journal of Bioscience and Bioengineering, 2010

Publication

1%

3

Fernanda G. Santos, Letícia C. Bonkovoski, Francielle P. Garcia, Thelma S. P. Cellet et al. "Antibacterial Performance of a PCL–PDMAEMA Blend Nanofiber-Based Scaffold Enhanced with Immobilized Silver Nanoparticles", ACS Applied Materials & Interfaces, 2017

1%

4

Khan, A.L.. "Ameliorative symbiosis of endophyte (*Penicillium funiculosum* LHL06) under salt stress elevated plant growth of *Glycine max* L.", *Plant Physiology and Biochemistry*, 201108

1%

Publication

5

Vigentini, I., G. De Lorenzis, V. Fabrizio, F. Valdetara, M. Faccincani, C. A. Panont, C. Picozzi, S. A. Imazio, O. Failla, and R. Foschino. "The vintage effect overcomes the terroir effect: a three years survey on the wine yeast biodiversity in Franciacorta and Oltrepo Pavese, two Northern Italian vine-growing areas", *Microbiology*, 2014.

1%

Publication

6

Nguyen, Long N. Ma, Dongliang Shui, Guanghou Wong, Peiyan Cazenave-Gassiot, Amaury Zhang, Xiaodong W. "Mfsd2a is a transporter for the essential omega-3 fatty acid docosahexaenoic acid.(LETTER)(Report)", *Nature*, May 22 2014 Issue

<1%

Publication

7

Nyaku, Seloame T. Sripathi, Venkateswara R. Kantety, Ramesh V. Cseke, Sarah B. Buyyarapu, Ramesh Mc . "Characterization of the reniform nematode genome by shotgun

<1%

8

Haziel Diana Jenifer, Sarthak Bhola, Veena Kalburgi, Shivaraj Warad, Vijayalaxmi M. Kokatnur. "The influence of cigarette smoking on blood and salivary super oxide dismutase enzyme levels among smokers and nonsmokers —A cross sectional study", Journal of Traditional and Complementary Medicine, 2015

Publication

<1 %

9

Pradeep, F. Stanly, M. Palaniswamy, S. Ravi, A. Thangamani, and B.V. Pradeep. "Larvicidal activity of a novel isoquinoline type pigment from Fusarium moniliforme KUMBF1201 against Aedes aegypti and Anopheles stephensi", Process Biochemistry, 2015.

Publication

<1 %

10

Avalos, Javier, and M. Carmen Limón. "Biological roles of fungal carotenoids", Current Genetics, 2015.

Publication

<1 %

11

Ungureanu, Camelia, Simona Popescu, Gabriela Purcel, Vlad Tofan, Marian Popescu, Aurora Sălăgeanu, and Cristian Pîrvu. "Improved antibacterial behavior of titanium surface with torularhodin–polypyrrole film",

<1 %

12

Lai Keng Chan. "Establishment of *Cyperus aromaticus* cell suspension cultures for the production of Juvenile hormone III", *In Vitro Cellular & Developmental Biology - Plant*, 09/16/2009

Publication

<1 %

13

T. Oda. "CEL-I, an Invertebrate N-Acetylgalactosamine-specific C-Type Lectin, Induces TNF- and G-CSF Production by Mouse Macrophage Cell Line RAW264.7 Cells", *Journal of Biochemistry*, 09/10/2007

Publication

<1 %

14

"*Catharanthus roseus*", Springer Nature, 2017

Publication

<1 %

15

Kajal Chakraborty, Bini Thilakan, Vamshi Krishna Raola. "Previously Undescribed Antibacterial Polyketides from Heterotrophic *Bacillus amyloliquefaciens* Associated with Seaweed *Padina gymnospora*", *Applied Biochemistry and Biotechnology*, 2017

Publication

<1 %

16

D.L Bishop. "Changes in carbohydrate coordinated partitioning and cell wall remodeling with stress-induced pathogenesis in wheat sheaths", *Physiological and Molecular Plant*

<1 %

17

Sandra R. Pombeiro-Sponchiado, Gabriela S. Sousa, Jazmina C.R. Andrade, Helen F. Lisboa, Rita C.R. Gonçalves. "Chapter 4 Production of Melanin Pigment by Fungi and Its Biotechnological Applications", InTech, 2017

Publication

<1 %

18

Rolf Henrik Nilsson. "The ITS region as a target for characterization of fungal communities using emerging sequencing technologies", FEMS Microbiology Letters, 07/2009

Publication

<1 %

19

Khudhair, Ameer Badr, Tony Hadibarata, Abdull Rahim Mohd Yusoff, Zee Chuang Teh, Liyana Amalina Adnan, and Hesam Kamyab. "Pyrene Metabolism by New Species Isolated from Soil Rhizoctonia Zeae SOL3", Water Air & Soil Pollution, 2015.

Publication

<1 %

20

Christiana, N Ogbonna. "Production of food colourants by filamentous fungi", African Journal of Microbiology Research, 2016.

Publication

<1 %

21

Ma, W.K.. "A PCR-DGGE method for detecting arbuscular mycorrhizal fungi in cultivated soils", Soil Biology and Biochemistry, 200509

<1 %

22

"ISOLATION IDENTIFICATION AND ENZYME PRODUCTION PROFILE OF A. NIGER.",
Journal of Animal and Plant Sciences, Oct 31
2014 Issue

<1 %

Publication

23

Wu, Bin, Vanessa Oesker, Jutta Wiese, Rolf Schmaljohann, and Johannes Imhoff. "Two New Antibiotic Pyridones Produced by a Marine Fungus, Trichoderma sp. Strain MF106", Marine Drugs, 2014.

<1 %

Publication

24

Claira Arul Aruldass, Laurent Dufossé, Wan Azlina Ahmad. "Current perspective of yellowish-orange pigments from microorganisms- a review", Journal of Cleaner Production, 2018

<1 %

Publication

25

"Posters", Clinical Microbiology and Infection, 2011

<1 %

Publication

26

Jin, Liming, Chunshan Quan, Xiyan Hou, and Shengdi Fan. "Potential Pharmacological Resources: Natural Bioactive Compounds from Marine-Derived Fungi", Marine Drugs, 2016.

<1 %

Publication

"Diversity and Benefits of Microorganisms from

27	the Tropics", Springer Nature, 2017 Publication	<1 %
28	Microbial Inoculants in Sustainable Agricultural Productivity, 2016. Publication	<1 %
29	Llenado, R. A., and G. A. Rechnitz. "Improved enzyme electrode for amygdalin", Analytical Chemistry, 1971. Publication	<1 %
30	ROMAN MARECIK, LIDIA BŁASZCZYK, RÓŻA BIEGAŃSKA-MARECIK, AGNIESZKA PIOTROWSKA-CYPLIK. "Screening and Identification of Trichoderma Strains Isolated from Natural Habitats with Potential to Cellulose and Xylan Degrading Enzymes Production", Polish Journal of Microbiology, 2018 Publication	<1 %
31	Bhaskar Bhadra. "Pichia cecembensis sp. nov. isolated from a papaya fruit (Carica papaya L., Caricaceae)", FEMS Yeast Research, 6/2007 Publication	<1 %
32	Frasz, Samantha L. Walker, Allison K. Ns. "Distribution of the foliar fungal endophyte Phialocephala scopiformis and its toxin in the crown of ", Canadian Journal of Forest Research, Sept 2014 Issue Publication	<1 %

33

Pandey, Veena, Mohammad W. Ansari, Suresh Tula, Sandep Yadav, Ranjan K. Sahoo, Nandini Shukla, Gurdeep Bains, Shail Badal, Subhash Chandra, A. K. Gaur, Atul Kumar, Alok Shukla, J. Kumar, and Narendra Tuteja. "Dose-dependent response of *Trichoderma harzianum* in improving drought tolerance in rice genotypes", *Planta*, 2016.

Publication

<1 %

34

Terhonen, Eeva, Teresa Marco, Hui Sun, Risto Jalkanen, Risto Kasanen, Martti Vuorinen, and Fred Asiegbu. "The effect of latitude, season and needle-age on the mycota of Scots pine (*Pinus sylvestris*) in Finland", *Silva Fennica*, 2011.

Publication

<1 %

35

"Studies from N. Waybright and co-researchers in the area of bird flu epidemiology published. (Report)", *World Disease Weekly*, Jan 27 2009 Issue

Publication

<1 %

36

Farah Naz Talpur. "Analysis and Characterization of Anthocyanin from Phalsa (*Grewia asiatica*)", *MOJ Food Processing & Technology*, 2017

Publication

<1 %

37

F. Delgado-Vargas. "Natural Pigments:

Carotenoids, Anthocyanins, and Betalains --
Characteristics, Biosynthesis, Processing, and
Stability", Critical Reviews in Food Science and
Nutrition, 5/1/2000

<1 %

Publication

38

Skoneczny, Dominik, Michał Oskiera,
Magdalena Szczech, and Grzegorz
Bartoszewski. "Genetic diversity of *Trichoderma*
atroviride strains collected in Poland and
identification of loci useful in detection of within-
species diversity", *Folia Microbiologica*, 2015.

<1 %

Publication

39

Candice Brinkmann, Philip Kearns, Elizabeth
Evans-Illidge, D. Kurtböke. "Diversity and
Bioactivity of Marine Bacteria Associated with
the Sponges *Candidaspongia flabellata* and
Rhopaloeides odorabile from the Great Barrier
Reef in Australia", *Diversity*, 2017

<1 %

Publication

40

Alapati Kavitha, Handanahal S. Savithri.
"Biological Significance of Marine Actinobacteria
of East Coast of Andhra Pradesh, India",
Frontiers in Microbiology, 2017

<1 %

Publication

41

Eva García-López, Alberto Alcázar, Ana María
Moreno, Cristina Cid. "Color-Producing
Extremophiles", *Wiley*, 2017

<1 %

Publication

A. Vilorio-Matos. "ISOLATION AND IDENTIFICATION OF BETACYANIN FROM FRUITS OF *Opuntia boldinghii* Br. et R. BY HPTLC AISLAMIENTO E IDENTIFICACION DE BETACIANINA PROVENIENTE DE FRUTOS DE *Opuntia boldinghii* Br. et R. POR HPTLC AILLAMENTO E IDENTIFICACION DE BETACIANINA PROVINTE DE FRUTOS DE *Opuntia boldinghii* Br. et R. POR HPTLC", CyTA - Journal of Food, 12/01/2001

Publication

Exclude quotes Off

Exclude matches Off

Exclude bibliography On